

Clinical Research

Official Publication of the American Federation for Clinical Research

LETTER OF TRANSMITTAL

Title of Abstract Quantitation of Serum Erythropoietic Stimulatory and Inhibitory

Activities Using ^3H -Thymidine Uptake by K562 Cells

All abstracts are published whether or not selected for presentation. Proper placement of the abstract in CLINICAL RESEARCH is dependent upon providing the information requested in the space below.

Organization: AFCR ; Nat. med meeting Meeting Date: 1st May 1987
(Specify organization and date of meeting)

NATIONAL:

American Federation for Clinical Research
American Society for Clinical Investigation
Association of American Physicians
Other (give full name of organization)

REGIONAL:

Eastern: Eastern Section, AFCR
Eastern Region, The Society for Investigative Dermatology
Midwest: Midwest Section, AFCR
Central Society for Clinical Research
Midwest Society for Pediatric Research
Central Region, The Society for Investigative Dermatology
Central Rheumatism Association
Southern: Southern Section, AFCR
Southern Society for Clinical Investigation
Southern Society for Pediatric Research
Southern Region, The Society for Investigative Dermatology
Southern Section, SREPCIM
Western: Western Section, AFCR
Western Society for Clinical Investigation
Western Society for Pediatric Research
Western Region, The Society for Investigative Dermatology

If electronic transmission were available,
would you take advantage of this service?

☒ YES ☐ NO

HIGHEST DEGREE(S)

eg, B.A., M.S.

M.D., Ph.D.

MEMBERSHIP*

STUDENT**

Author's Name (age of nonmembers and presenter must be given)	AGE	HIGHEST DEGREE(S)	Active	Senior	Associate	Non	A	B
1. Lewis, J. P.	69	Ph.D.				X		
2. Felice, A. E.	38	M.D., Ph.D.	X					
3. _____								
4. _____								
5. _____								
6. _____								
Introducer (if applicable)								

*Check appropriate box(es) for each author.

**Check Box A if the student initiated, designed, and executed the research; check Box B if the student was a collaborator.

Paper (if selected) will be presented by A. E. Felice

whose age at time of meeting will be 39 years.

(name of author)

Members are reminded that the presenter may not be over age 41 unless there are no other younger coauthors (Bylaw Article VI Section 2)

☐ For publication only. This box should be checked ONLY if the abstract is not to be considered for presentation.

The sponsoring member affirms: (1) personal responsibility for assuring that at the time of the meeting for which it is being considered the material herein will not have been published or presented at any national meeting; (2) that if accepted, an eligible author is prepared to present it; (3) that any animal studies conform with the "Guiding Principles in the Care and Use of Animals" of the American Physiological Society; and (4) that any human experimentation was conducted according to a protocol approved by an Institutional Review Board on ethics of human investigation (if no such Board exists, please state in a separate letter what standards were used to assess the ethical propriety of the protocol).



Member's signature

Abstract Reproduction Form

AFCR

(name of organization, selected from list on form letter of transmittal)

TYPE name, address, and telephone number of author who should receive correspondence in Box A and complete Boxes B, C, and D.

Telephone (404) 724-5116, x2457 (404) 733-2646
(Area code) office (Area code) home

A

Name A. E. Felice, M.D., Ph.D.
Address Hemoglobin Research Laboratory, Medical Research Service, Veterans Administration Medical Center, Augusta, GA 30910

B (See Rule 5)

(A separate check must accompany each abstract)

Date _____

Payment (\$30.00) _____

Check # _____

Purchase order # (\$35.00) _____

Issued by _____

(name of institution)

A copy of this abstract must be attached to original purchase order to aid in identification.

C

Do not consider for poster session ☒

D

CHECK SINGLE SUBSPECIALTY CLASSIFICATION:

Allergy _____
Cardiovascular* _____ Code No. _____
Clinical Epidemiology—
Health Care Research _____
Clinical Nutrition _____
Clinical Pharmacology _____
Critical Care Medicine _____
Dermatology _____
Endocrinology (see Rule 6) _____
Gastroenterology _____
Genetics _____
Hematology _____ ☒
Hypertension _____
Immunology _____
Infectious Disease _____
Metabolism (see Rule 6) _____
Oncology _____
Pulmonary _____
Renal & Electrolyte _____
Rheumatology _____

*For abstracts submitted to cardiovascular only, select single subcategory and enter code no. (1-5) in space above: (1) Cardiac biochemistry/cell biology; (2) Hemodynamics/reflexes; (3) Electrophysiology/dysrhythmias; (4) Coronary atherosclerosis/lipids; (5) Noninvasive techniques. Subclassifications is designed to aid in reviewing process only and is independent of program selection.

FEATURED RESEARCH SYMPOSIA
(for National Meeting only)

Atrial Natriuretic Factors _____
Endothelial Cells/Atherosclerosis _____
Growth Factors and Cellular Proliferation _____
Guanine-Nucleotide Regulatory Proteins _____
Receptors _____
Regulation of Gene Expression _____
Regulation of T-Cell Function _____
Restriction Fragment Length Polymorphism _____
Therapy of AIDS _____
Thrombolytic Therapy in Cardiovascular Disease _____

QUANTITATION OF SERUM ERYTHROPOIETIC STIMULATORY AND INHIBITORY ACTIVITIES USING ^3H -THYMIDINE UPTAKE BY K562 CELLS. J. P. Lewis* and A. E. Felice**, Hemoglobin Research Laboratory, Medical Research Service, Veterans Administration Medical Center, and Departments of Cell and Molecular Biology, Pediatrics, and Medicine, Medical College of Georgia, Augusta, GA.

Cells of the human pluripotent leukemia cell line K562 exhibit erythroid properties such as glycophorin and hemoglobin synthesis. We examined the response of K562 cells to erythropoietin in terms of ^3H -thymidine uptake as described by Krystal et al for spleen cells from anemic mice pretreated with phenylhydrazine (Exp. Hematol. 11: 649, 1983). Both murine spleen cells and K562 cells gave the same dose response curve, i.e. 10^3 cpm/ 10^3 cells/mU erythropoietin. Thus, the K562 cells could be used instead of the murine spleen cells in the Krystal assay for erythropoietic activity in serum. With this assay, nine healthy volunteers with normal blood counts gave erythropoietin levels of 356 ± 10 (SD) mU/ml. Sera of 24 children with sickle cell disease gave levels between 289 and 1835 mU/ml (average is 589 ± 66 (SD) mU/ml). Addition of an anti-erythropoietic inhibitory factor antiserum, or dilution of sera led to increased ^3H -thymidine uptake. Thus, ^3H -thymidine uptake reveals the presence of physiological erythropoietin receptors on K562 cells and can be used to measure erythropoietic and anti-erythropoietic activities in sera of patients.

BOTH THIS FORM AND THE FORM LETTER OF TRANSMITTAL MUST BE SIGNED BY A MEMBER (RULE 2)

A. E. Felice

(please type name)

MEMBER'S SIGNATURE _____

Revised October 1986